arXiv:1204.5300v3 [cond-mat.str-el] 1 May 2012

Revival of Single-Particle Transport Theory for the Normal State of High-T_c Superconductors: I. Relaxation-Time Approximation

O. Narikiyo *

(Apr. 24, 2012)

Abstract

A demonstration of the usefulness of a simple formula for the DC Hall conductivity based on the relaxation-time approximation is presented. The implementation of the formula on your PC can be completed within ten minutes. The formula is employed to explain the qualitative features of the measured DC Hall conductivity in the normal state of high-T_c superconductors.

A revival of a single-particle transport theory for the normal state of high-Tc superconductors is now in progress [1, 2, 3].

In this Short Note I try to support the progress using a simple formula for the DC Hall conductivity. The formula is so simple that it can be implemented on your PC within ten minutes.

The target data is the non-monotonic temperature dependence of the DC Hall conductivity measured in the normal state of PCCO superconductors [4]. Although it was interpreted [4] by the effect of the vertex correction, I shall show that it is explained qualitatively within the relaxation-time approximation.

In the next Short Note I shall discuss why the relaxation-time approximation is enough for the qualitative understanding and the vertex correction is irrelevant.

*Department of Physics, Kyushu University, Fukuoka 812-8581, Japan

We start from the full Green function for electrons

$$G^R(\mathbf{p},\epsilon) = \frac{1}{\epsilon - E(\mathbf{p},\epsilon) + i/2\tau(\mathbf{p},\epsilon)}, \qquad G^A(\mathbf{p},\epsilon) = \frac{1}{\epsilon - E(\mathbf{p},\epsilon) - i/2\tau(\mathbf{p},\epsilon)}, \tag{1}$$

where the self-energy is renormalized into the dispersion $E(\mathbf{p},\epsilon)$ and the life-time $\tau(\mathbf{p},\epsilon)$.

Our numerical calculation is done for 2D square lattice with the lattice constant $a \equiv 1$. By the symmetry we only need the information for the quarter of the Brillouin zone: $0 < p_x < \pi$ and $0 < p_y < \pi$. In the following the summation over $\mathbf{p}$ is restricted within this quarter.

We consider the conductivity within the relaxation-time approximation. In the next Short Note we shall discuss that the vertex correction does not change the qualitative result of the relaxation-time approximation in the case where the forward scattering is unimportant. Neglecting the vertex correction the DC conductivity σ_{xx} in the absence of the magnetic field is given by [5]

$$\sigma_{xx} = \frac{e^2}{\pi} \cdot \frac{4}{R(T)}, \tag{2}$$

with

$$\frac{1}{R(T)} \equiv \sum_{\mathbf{p}} v_x^2 G^R G^A = \sum_{\mathbf{p}} v_x^2 \frac{1}{E^2 + (1/2\tau)^2}, \tag{3}$$

where $G^R = G^R(\mathbf{p},0)$, $G^A = G^A(\mathbf{p},0)$, $E = E(\mathbf{p},0)$, $\tau = \tau(\mathbf{p},0)$ and $v_x = \partial E/\partial p_x$. Here we have assumed the Fermi degeneracy for simplicity.

In the same manner the DC Hall conductivity σ_{xy} proportional to the weak magnetic field H is given by [5]

$$\sigma_{xy}/H = \frac{|e|^3}{\pi} \cdot 4S(T), \tag{4}$$

with

$$\begin{aligned} S(T) &\equiv \sum_{\mathbf{p}} v_x \Big(v_x \frac{\partial v_y}{\partial p_y} - v_y \frac{\partial v_x}{\partial p_y} \Big) G^R G^A \frac{G^R - G^A}{2i} \\ &= -\sum_{\mathbf{p}} v_x \Big(v_x \frac{\partial v_y}{\partial p_y} - v_y \frac{\partial v_x}{\partial p_y} \Big) \frac{1/2\tau}{\big[E^2 + (1/2\tau)^2\big]^2}, \end{aligned} \tag{5}$$

where $v_y = \partial E/\partial p_y$.

Thus, if you have the data of the momentum dependence of the dispersion E and the life-time τ at $\epsilon = 0$, you can easily get the DC Hall conductivity from this formula. The implementation of the formula on your PC takes little effort. I employ the following model for E and τ.

Here we substitute the dispersion obtained by the band calculation [6]

$$E = -2t(\cos p_x + \cos p_y) + 4t' \cos p_x \cdot \cos p_y - 2t''(\cos 2p_x + \cos 2p_y) - \mu, \quad (6)$$

for that for quasi-particles. For PCCO we adopt $t = -0.438$eV, $t' = 0.156$eV and $t'' = 0.098$eV [6] in the numerical calculations in this Short Note. By this substitution we miss some many-body correlations.

The Fermi surface for this dispersion is shown in Fig. 1. The factor $f(\mathbf{p}) \equiv v_x(v_x \partial v_y/\partial p_y - v_y \partial v_x/\partial p_y)$ appearing in σ_{xy} is shown in Fig. 2.

For the life-time we adopt the model

$$\frac{1}{2\tau} = w(\mathbf{p}) \cdot g_1 \cdot T + \big[1 - w(\mathbf{p})\big] \cdot g_2 \cdot T^2, \quad (7)$$

similar to the multi-patch model [7] where the Brillouin zone is divided into hot and cold patches. The life-time is determined by the coupling to the anti-ferromagnetic spin-fluctuation. In the cold patches the relevant spin-fluctuation spectrum is broad and its temperature-dependence is weak so that $1/\tau \sim T^2$. On the other hand, in the hot patches it is peaked around the nesting-vector and its integrated weight depends on the temperature so that $1/\tau \sim T$. For simplicity we did not implement the gradual change between hot and cold patches employed in [7]. Namely $w(\mathbf{p})$ is a step function in our case where $w(\mathbf{p}) = 1$ if $(\mathbf{p} - \mathbf{p}_1)^2 < r^2$ or $(\mathbf{p} - \mathbf{p}_2)^2 < r^2$ with $\mathbf{p}_1 \equiv (\pi, 0)$ and $\mathbf{p}_2 \equiv (0, \pi)$ and $w(\mathbf{p}) = 0$ otherwise.

We set $g_2 = 100/\text{eV}$ in accordance with [3]. Here the temperature T is measured in eV. On the other hand, our choice, $g_1 = 0.1$, is smaller than [3] and [7]. Such a smallness is explained by the nested spin fluctuation [8].

Before performing the numerical calculation we can make a rosy prediction that the non-monotonic temperature dependence of σ_{xy}, which we want to derive, is obtained, if the temperature dependences of $1/\tau$ are different between the regions with positive $f(\mathbf{p})$ and the regions with negative $f(\mathbf{p})$ in the Brillouin zone. After the numerical calculation we confirm the prediction as shown in Fig. 3. The result for $\mu = 0$ actually shows the non-monotonic temperature dependence qualitatively similar to that observed in the experiment [4]. More direct comparison between the experiment [4] and our numerical calculation can be done for σ_{xy}/σ_{xx} shown in Fig. 4.

In conclusion our simple formula based on the relaxation-time approximation for single particles qualitatively explain the non-monotonic behavior of the DC Hall conductivity in the normal state of PCCO superconductors, though our choice of the parameters should be reconsidered for quantitative description.

This work was driven by the discussions with Professor Kazumasa Miyake.

References

[1] P. A. Casey and P. W. Anderson: Phys. Rev. Lett. **106**, 097002 (2011).

[2] J. Kokalj and R. H. McKenzie: Phys. Rev. Lett. **107**, 147001 (2011).

[3] J. Kokalj, N. E. Hussey and R. H. McKenzie: arXiv:1202.4820.

[4] G. S. Jenkins, D. C. Schmadel, P. L. Bach, R. L. Greene, X. Béchamp-Laganière, G. Roberge, P. Fournier, H. Kontani, and H. D. Drew: Phys. Rev. B **81**, 024508 (2010).

[5] H. Fukuyama, H. Ebisawa, and Y. Wada: Prog. Thoer. Phys. **42**, 494 (1969).

[6] I. A. Nekrasov, E. Z. Kuchinskii, and M. V. Sadovskii: J. Phys. Chem. Solids **72**, 371 (2011).

[7] A. Perali, M. Sindel, and G. Kotliar: Eur. Phys. J. B **24**, 487 (2001).

[8] O. Narikiyo and K. Miyake: Physica C **307**, 254 (1998).



Figure 1: Fermi surfaces for $\mu = -0.1$, 0, 0.1 with $t = -0.438$, $t' = 0.156$, $t'' = 0.098$ where all the energies are represented in eV. Only the quarter of the Brillouin zone ($0 < p_x < \pi$ and $0 < p_y < \pi$) is depicted.



Figure 2: The factor $f(\mathbf{p}) = v_x(v_x\partial v_y/\partial p_y - v_y\partial v_x/\partial p_y)$ in the quarter of the Brillouin zone. In the white region $f(\mathbf{p}) > 0$ and $f(\mathbf{p}) < 0$ in the black region.



Figure 3: Temperature dependencies of $S(T)$ for $\mu = -0.1,\ 0,\ 0.1$ with $r^2 = 1.07$ where $S(T) \propto \sigma_{xy}$. The temperature is represented in eV. The summation over $\mathbf{p}$ is carried out using 3000×3000 mesh in the quarter of the Brillouin zone.



Figure 4: The temperature dependence of $S(T)R(T)$ for $\mu = 0$ with $r^2 = 1.07$ where $S(T)R(T) \propto \sigma_{xy}/\sigma_{xx}$.